

References

1. Fried, E. & Nesse, R.M. (2015) Depression is not a consistent syndrome: an investigation of unique symptom patterns in the STAR*D study. *J Affect Disord*, 172, 96–102.
2. Beck, A. (1967) *Depression: Clinical, Experimental, and Theoretical Aspects*. Harper and Row, New York.
3. Dean, J. & Keshavan, M. (2017) The neurobiology of depression: An integrated view. *Asian J. Psychiatry*, 27, 101–111.
4. Disner, S.G., Beevers, C.G., Haigh, E.A., & Beck, A. (2011) Neural mechanisms of the cognitive model of depression. *Nat Rev Neurosci*, 12, 467–477.
5. Kraus, C., Kadriu, B., Lanzenberger, R., Zarate, C.A., & Kasper, S. (2019) Prognosis and improved outcomes in major depression: a review. *Transl. Psychiatry*, 9, 127.
6. Ruhé, H.G., van Rooijen, G., Spijker, J., Peeters, F.P.M.L., & Schene, A.H. (2012) Staging methods for treatment resistant depression. A systematic review. *J. Affect. Disord.*, 137, 35–45.
7. Bewernick, B.H., Kayser, S., Gippert, S.M., Switala, C., Coenen, V.A., & Schlaepfer, T.E. (2017) Deep brain stimulation to the medial forebrain bundle for depression- long-term outcomes and a novel data analysis strategy. *Brain Stimulat.*, 10, 664–671.
8. Drobisz, D. & Damborská, A. (2019) Deep brain stimulation targets for treating depression. *Behav. Brain Res.*, 359, 266–273.
9. Williams, L.M. (2016) Precision psychiatry: a neural circuit taxonomy for depression and anxiety. *Lancet Psychiatry*, 3, 472–480.
10. Li, B.-J., Friston, K., Mody, M., Wang, H.-N., Lu, H.-B., and Hu, D.-W. (2018). A brain network model for depression: From symptom understanding to disease intervention. *CNS Neurosci. Ther.* 24, 1004–1019. doi: 10.1111/cns.12998.
11. Coenen, V. A. et al. Tractographic description of major subcortical projection pathways passing the anterior limb of the internal capsule. Corticopetal organization of networks relevant for psychiatric disorders. *Neuroimage Clin* 25, 102165 (2020).